

Consistency of Wilsonian effective actions

Vipul Periwal

Department of Physics, Princeton University, Princeton, New Jersey 08544

Wilsonian effective actions are interpreted as free energies in ensembles with prescribed field expectation values and prescribed connected two-point functions. Since such free energies are directly obtained from two-particle-irreducible functionals, it follows that Wilsonian effective actions satisfy elementary perturbative consistency conditions, and non-perturbative convexity conditions. In particular, the exact determination of a Wilsonian action by other means (e.g. supersymmetry) allows one to extract restrictions on the particular cutoff scheme and field reparametrization that would lead to such a Wilsonian action from an underlying microscopic action.

The Wilsonian effective action [1] figures prominently in the spectacular advances [2,3] that have been made in the understanding of the dynamical properties of supersymmetric quantum field theories. This kind of effective action does not involve integrating out massless degrees of freedom, hence has locality properties that allow one to utilize properties of supersymmetric actions such as holomorphy that do not hold for the 1PI effective action when massless particles are present. Of course, the Wilsonian action is to be used in a functional integral over the remaining long wavelength modes, unlike the 1PI action. Based on such properties, remarkable arguments have been given to determine the low-energy dynamics of supersymmetric field theories [4]. There appears to be great predictive power in this approach.

Given the physical importance of the Wilsonian effective action, it is natural to ask if there is a regulator-independent physical characterization of the Wilsonian action, analogous to the interpretation of the 1PI effective action as the free energy in an ensemble with a prescribed field expectation value. This question becomes especially interesting in light of the unease that has been expressed [5] regarding the holomorphy assumptions that play a large part in the supersymmetric applications mentioned above. Ref. [5] argued, following Shifman and Vainshtein [6], that not just the microscopic action, but also the exact ‘field-dependent cutoff’ scheme and possible field reparametrizations, must be regarded as additional parameters in low-energy Wilsonian actions. This would seem to open up a veritable Pandora’s box of arbitrariness in deductions based on Wilsonian effective actions.

It would appear to follow then that the exact determination of a low-energy Wilsonian action should allow the determination of the cutoff scheme, and the field reparametrization, that leads from the microscopic action to the Wilsonian action. In particular, if one has deduced the exact non-derivative terms in the Wilsonian action, one might be able to construct (or at least find restrictions on the form of) the rest of the Wilsonian action from such knowledge. This is, potentially, of great interest for a variety of reasons (see below). Unfortunately, one cannot carry this program out for lack of a scheme-independent characterization of the Wilsonian action.

The aim of this note is to present a simple and explicit scheme for extracting such information, by realizing that an alternative physical characterization of the Wilsonian action can be given in terms of the free energy in a particular ensemble. This physical interpretation is non-perturbative, and makes no explicit reference to any cutoff scheme. It therefore places the Wilsonian action in a context that allows one to ‘reverse engineer’ the cutoff scheme from the Wilsonian action. In fact, I shall show that the Wilsonian action is directly obtained from the two-particle-irreducible(2PI) functional. Since the 2PI functional has a particular perturbative expansion, one can match the Wilsonian action with this perturbative expansion, and hence restrict the cutoff scheme and field reparametrization that led to the Wilsonian action. For example, if presented with an action and the information that it is a Wilsonian effective action, it is possible to test such assertions in a straightforward manner. Quite apart from these perturbative consistency conditions, the free energy interpretation of the Wilsonian action implies non-perturbative convexity properties, analogous to the usual convexity of the 1PI effective action. In essence, the 2PI formalism enables one to phrase cutoff dependence directly in terms of the properties of the non-perturbative connected two-point function.

The organization of this paper is as follows: I first review some aspects of the 2PI functional, following Cornwall, Jackiw and Tomboulis [7]. I then show that the Wilsonian effective action is the free energy in an ensemble with prescribed field expectation value and *prescribed* connected two-point correlation function. Since such a free energy is just a particular case of the 2PI functional, an infinite number of perturbative consistency checks follow immediately from the properties of the 2PI functional. I show in some examples how the cutoff is restricted in this interpretation. Some comments on the utility of this interpretation are in a concluding paragraph.

The 2PI functional, in the form I need, was derived by Cornwall, Jackiw and Tomboulis [7] in an unrelated context; it was initially studied in by Lee and Yang [8], and others [9]. It has been used in the study of the bound state problem in quantum field theory, where it can be used to extract non-perturbative information via variational principles. I give a short account of the definition of the 2PI functional, with some useful properties, in Euclidean space, for convenience. This is essentially taken from [7], replacing various factors of i .

If S_0 is a microscopic action, define a generating functional

$$W[J, K] \equiv \ln \int D\phi \exp(-S_0[\phi] + J\phi + \frac{1}{2}\phi K\phi),$$

where $J\phi \equiv \int dx J(x)\phi(x)$, and $K(x, y)$ is a source for the complete two-point function. Define the Legendre transform of W

$$\Gamma[\varphi, G] + W[J, K] \equiv J\varphi + \frac{1}{2}(\varphi K\varphi + GK), \quad (1)$$

($GK \equiv \int dxdy G(x, y)K(x, y)$) which implies that

$$\frac{\delta\Gamma}{\delta\varphi} = J + K\varphi, \quad \frac{\delta\Gamma}{\delta G} = K. \quad (2)$$

Since

$$\frac{1}{2}\langle\phi\phi\rangle = \frac{\delta W}{\delta K} = \frac{1}{2}(G + \varphi\varphi),$$

it follows that G is the exact connected two-point function, and $\varphi = \langle\phi\rangle$ as usual.

Γ is given, in perturbation theory, by a sum of graphs that have G as the propagator in internal lines, and $\delta^n S_0[\varphi]/\delta\varphi^n$ vertices ($n > 2$). Since G is the exact two-point function, there are no self-energy corrections in this set of graphs; in other words, these graphs are all 2PI. There are just two contributions at one-loop order, $\frac{1}{2} \int dxdy G(x, y) \frac{\delta^2 S_0[\varphi]}{\delta\varphi(x)\delta\varphi(y)}$, and $-\frac{1}{2}\text{Tr} \ln G$ (apart from a term which is G and φ independent). The formal derivation is given in [7].

What does all this have to do with the Wilsonian effective action? Recall that Polchinski's derivation of the Wilsonian renormalization group equation [10] involved the addition of a cutoff function to the bare action. The cutoff is introduced by adding a term of the form $\int \frac{1}{2}\phi K\phi$ to the action, where the Fourier transform of K is taken to vanish for $p^2 < \Lambda^2$, and $-K/p^2$ tends to infinity rapidly for $p^2 > \Lambda^2$. (The function K used here is different from K in Polchinski's paper: $K_{\text{Polchinski}}(p^2/\Lambda^2) \equiv (1 - K/p^2)^{-1}$.) Thus $W[J, K]$ contains the information needed to implement the Wilsonian effective action, for appropriate choices of K .

However, K is an inherently perturbative object—thinking in terms of K requires thinking in terms of Feynman diagrams. We should focus instead on physical quantities, such as G ; the awkward description of K tending to infinity is expressed as the vanishing of G for the appropriate set of modes. Formally, from Eq. 2, and the usual relation

$$\Gamma[\varphi] + W[J] = \varphi J \quad (3)$$

we note that modes for which K and J vanish have, in fact, been integrated out—the partition function has been exactly evaluated as an integral over these modes. On the other hand, modes for which G vanishes, are uncorrelated in a non-perturbative sense. It is trivial to see that the perturbative description of this non-perturbative characterization is that there are no internal lines involving the propagation of such modes in Feynman diagrams.

To see that this is precisely what the Wilsonian effective action requires, let us suppose that we have two fields ϕ, Φ , and we want the Wilsonian effective action for ϕ obtained by integrating out Φ . Then the claim is that

$$S_W(\phi) = \Gamma[\phi, \Phi_s, G_\phi = 0, G_{\Phi, s}]$$

where

$$\Phi_s : \frac{\delta\Gamma}{\delta\Phi_s} = 0, \quad G_{\Phi, s} : \frac{\delta\Gamma}{\delta G_{\Phi, s}} = 0. \quad (4)$$

Clearly, Eq. 4 implies that the integral over Φ has been carried out exactly, and $G_\phi = 0$ is precisely the non-perturbative analogue of the statement that there are no internal lines with ϕ contractions. Note that, in general, $\Phi_s = \Phi_s(\phi)$, $G_{\Phi, s} = G_{\Phi, s}(\phi)$.

This identification is pretty much content-free if one could use hard cutoffs, with no field dependence. Unfortunately, for the applications of interest, one cannot use such simple cutoffs. G *cannot* be taken to vanish abruptly, in which case one must find the appropriate smooth transition from $G = 0$ to $G : \frac{\delta\Gamma}{\delta G} = 0$, which preserves desirable properties such as holomorphy. The point is that G is an object with a non-perturbative meaning—it is the connected two-point function in the ensemble of interest. The field dependence of cutoffs can be stated in a precise manner now, as φ -dependence of the transition from $\frac{\delta\Gamma[\varphi, G]}{\delta G} = 0$ to $G = 0$.

As a first example, suppose that we are given a low-energy effective action for a real scalar field at scale Λ_R in four dimensions and told that the mass parameter has no quadratic dependence on Λ_R . For a general microscopic action

$$S_0[\phi] = \int \frac{1}{2} \phi(p^2 + m^2) \phi + \frac{g}{4!} \phi^4$$

with microscopic cutoff Λ held fixed, this implies that

$$Z_m m^2(\Lambda_R) = m^2 + g \int^\Lambda d^4 p G(p^2)$$

where $G = 0$ for $p^2 < \Lambda_R^2$. For $m^2(\Lambda_R)$ to have no quadratic dependence on Λ_R , we must have

$$\tilde{G}(x) \equiv xG(x) : \int_0^{(\Lambda/\Lambda_R)^2} dx x \tilde{G}' = 0.$$

(A similar equation appears in an appendix (eq. A.17) in [11], but these authors did not make the following observations.) In other words, the cutoff must be such that the correlation between modes being integrated out does not monotonically decay—it *must* oscillate, so that $\tilde{G}'$ is not a positive definite function. It is difficult to see how such a cutoff could be consistent with unitarity *given* the assumed microscopic dynamics, which involves no other degrees of freedom. Thus, in this example, we find that the assumed form of the low-energy Wilsonian action suggests the existence of other degrees of freedom. This is reminiscent of Intriligator’s arguments for ‘integrating in’ degrees of freedom [12].

As a second example, consider an $N = 1$ supersymmetric model in four dimensions, with just chiral multiplets. In this case, the interaction terms in a renormalizable microscopic superspace action are purely cubic and all two-loop and higher contributions to the Wilsonian action depend only on $G_{\Phi\Phi}$, $G_{\Phi\bar{\Phi}}$, and $G_{\bar{\Phi}\bar{\Phi}}$. It is then simple to see that any transition to $G_{\Phi\Phi} = 0$ that does not violate the tree-level θ dependence will imply the standard superpotential non-renormalization theorem [13].

In conclusion, the characterization of the Wilsonian effective action given in this note should be useful in restricting higher derivative terms, which is of some importance in, *e.g.* D3-brane scattering [14]. It should also be useful for figuring out potentially interesting field-dependent regulators for non-supersymmetric situations. As shown in the first example, it may be important for ruling out certain regulators as unphysical. The free energy interpretation implies *non-perturbative* convexity properties for the Wilsonian action, to which I hope to return elsewhere.

I am grateful to S. Minwalla, M. Roček and R. von Unge for helpful conversations. This work was supported in part by NSF grant PHY96-00258.

-
- [1] K.G. Wilson and J.B. Kogut, Phys. Rep. 12C 75 (1974)
 - [2] N. Seiberg and E. Witten, Nucl. Phys. B426, 19 (1994); B431, 484 (1994)
 - [3] N. Seiberg, Phys. Lett. B318, 469 (1993); Nucl. Phys. B435, 129 (1995); K. Intriligator and N. Seiberg, Nucl. Phys. Proc. Supp. 45BC, 1 (1996)
 - [4] For reviews, see M. Peskin, hep-th/9702094; L. Alvarez-Gaumé and F. Zamora, hep-th/9709180
 - [5] M. Dine and Y. Shirman, Phys. Rev. D50, 1362 (1994); S. de Alwis, hep-th/9508053; F. Bastianelli, Phys. Lett. B369, 249 (1996)
 - [6] M. Shifman and A. Vainshtein, Nucl. Phys. B359, 571 (1991)
 - [7] J.M. Cornwall, R. Jackiw and E. Tomboulis, Phys. Rev. D10, 2428 (1974)
 - [8] T.D. Lee and C.N. Yang, Phys. Rev. 117, 22 (1960)
 - [9] J.M. Luttinger and J.C. Ward, Phys. Rev. 118, 1417 (1960); P. Martin and C. de Dominicis, J. Math. Phys. 5, 14 (1964); H.D. Dahmen and G. Jona-Lasinio, Nuovo Cim. 52A, 807 (1967); A.N. Vasil’ev and A.K. Kazanskii, Theor. Math. Phys. 12, 875 (1972)
 - [10] J. Polchinski, Nucl. Phys. B231, 269 (1984)
 - [11] J. Hughes and J. Liu, Nucl. Phys. B307, 183 (1988)
 - [12] K. Intriligator, Phys. Lett. B336, 409 (1994)
 - [13] S.J. Gates, M.T. Grisaru, M. Roček and W. Siegel, *Superspace*, (Benjamin-Cummings, Reading MA, 1983)
 - [14] V. Periwal and R. von Unge, hep-th/9801121